



# Raw diets

Raw diets (also known as raw-meat-based diets, 'biologically appropriate raw food' or 'BARF' diets) are becoming increasingly popular with owners and, consequently, veterinary professionals need to provide guidance about the advantages and disadvantages of such foods.

## What are raw diets?

Raw diets predominately comprise uncooked ingredients, although different components can vary. Raw diets may contain:

- Meat
- Fish
- Poultry
- Bones
- Meat by-products (offal)
- Unpasteurized milk
- Uncooked egg
- Fruit and vegetables
- Nuts and seeds
- Oils
- Cereals
- Supplements.

## Types of raw diet

- Home-prepared:
  - Feeding to a recipe
  - No recipe; any available food is fed.
- Prey-model – whole animal carcasses are fed.
- Commercially manufactured – within this category, various types are offered, including:
  - Fresh
  - Frozen
  - Freeze-dried
  - Carbohydrate premix with a raw protein source
  - Kibble with freeze-dried raw pieces
  - Diets with a raw-meat coating.


**NB:** Recipes may not be complete and balanced. Commercially prepared raw diet recipes should be created by professional formulators trained in the commercial formulation of pet foods. Ideally, the formulation should also have been developed in consultation with an appropriately qualified individual.

- **Appropriate qualifications include:**
  - **PhD in animal nutrition**
  - **Diplomate of the American College of Veterinary Nutrition (ACVN)**
  - **Diplomate of the European College of Veterinary Comparative Nutrition (ECVCN).**

In the case of home-prepared raw diets, it is recommended that the recipe be designed, overseen or approved by a Diplomate of either the ACVN or the ECVCN.

## What are the advantages of feeding raw diets?

### Evidence-based advantages of raw diets

- Currently, there is insufficient published scientific evidence confirming many of the suggested benefits of raw diets. However, there is some emerging evidence for the following:
  - Dental benefits from chewing raw bones with meat: 

Delivered by BSAVA to:

- There is some evidence to suggest that the texture of chewing raw bones with meat could be beneficial for dental calculus. However, there is no evidence that it improves oral health or prevents plaque, periodontitis or tooth loss<sup>1</sup>. Bones, raw or cooked, can also pose a risk of dental fractures, oesophageal obstructions and foreign bodies.
- Improvement in coat quality<sup>2</sup>:
  - It has been suggested that this might be the result of feeding a diet with an increased fat content. Of course, any benefits of feeding a high-fat diet should be balanced against potential risks (e.g. increased risk of obesity and pancreatitis).
- Improved digestibility:
  - This is partially true in that diets based on raw meat are typically more digestible than dry proprietary food, but not necessarily wet proprietary food<sup>3</sup>
  - Digestibility varies depending on the nutrient<sup>4</sup>.
- Improved palatability<sup>5</sup>:
  - Food with a greater moisture content will be more palatable to some dogs and high protein is more palatable for cats. This may be an advantage for picky eaters, but could be a disadvantage for pets prone to obesity.

Proponents of raw diets also report the following advantages from their clinical experience:

Diet claim	Available evidence
A 'natural' food is fed, and owners may feel that this is better for the pet	<ul style="list-style-type: none"> <li>■ Guidance for the term 'natural' is that it must be obtained exclusively (or at least 95%) from the source material (e.g. of vegetable or animal origin). This is not an official or regulated definition<sup>5</sup></li> <li>■ Therefore, depending on the ingredients used, commercially manufactured diets can also be defined as 'natural' in accordance with the above guidance</li> </ul>
More appropriate for dogs. As wolves are similar to dogs, the sort of diet that they consume is thought to be more appropriate for dogs than processed kibble or canned food	<ul style="list-style-type: none"> <li>■ A study has shown that, over the many years of dog domestication, their dietary needs have changed and they have adapted their digestive capabilities to live as companions<sup>6</sup>. The typical lifespan of wolves is shorter than that of domesticated dogs and, although many factors are responsible for mortality in wolves and dogs, this evidence is not consistent with a significant benefit for raw food over proprietary food. Therefore, further work is required before any perceived advantages of a so-called 'ancestral diet' can be clarified</li> </ul>
Does not contain carbohydrate/gluten/wheat: <ul style="list-style-type: none"> <li>■ Carbohydrates are thought not to be digestible and may even be harmful to pets</li> <li>■ Carbohydrates are responsible for dietary intolerances</li> </ul>	<ul style="list-style-type: none"> <li>■ Unlike wolves, domesticated dogs can readily digest carbohydrates: indeed, 10 of the 36 genomic differences between these species are involved in carbohydrate digestion<sup>6</sup></li> <li>■ Studies have shown that cats can also utilize carbohydrates, but to a lesser extent than dogs<sup>7</sup></li> <li>■ True dietary intolerances are rare and generally assumed to be related to the protein component rather than carbohydrate components of a diet<sup>8,9,10</sup></li> </ul>
Reduced occurrence of gastrointestinal signs	<ul style="list-style-type: none"> <li>■ There is no published scientific evidence for this claim</li> </ul>
Improved protein digestibility	<ul style="list-style-type: none"> <li>■ True for some but not all proteins. If raw egg is given, the avidin in the egg white binds to the amino acid biotin, making the proteins in the diet unavailable and therefore much less digestible<sup>11</sup></li> </ul>
Improved stool volume due to feeding a diet without perceived 'fillers'	<ul style="list-style-type: none"> <li>■ No scientific study evidence</li> <li>■ The smaller stool volume in pets fed raw diets may be due to the lower fibre content<sup>3,12</sup></li> <li>■ There are no such things as 'fillers' in pet foods. Fillers are thought to be the low-quality ingredients listed as carbohydrate or starch sources in pet foods. They have also been described as 'cereal' or even as the 'sawdust content' of proprietary foods. It is thought that other than acting as a 'bulking' agent, they offer little or no nutritional benefit and, in some cases, may even cause harm<sup>13</sup>. It is also thought that they cause dilution of the nutrition within the diets and are present only to bring costs down<sup>14</sup>. There is no scientific evidence of this. There is, however, evidence that carbohydrate sources can deliver nutrients that can be easily utilized by dogs and cats and form part of a nutritious diet<sup>15</sup></li> </ul>
Improved stool quality	<ul style="list-style-type: none"> <li>■ No scientific study evidence, although stool volume is often less due to raw diets typically having a lower fibre content</li> </ul>
Improved breath odour	<ul style="list-style-type: none"> <li>■ No scientific study evidence</li> </ul>

Delivered by BSAVA to:

Diet claim	Available evidence
Reduced risk of obesity	<ul style="list-style-type: none"> <li>No scientific study evidence</li> <li>65% of dogs in the UK are either overweight or have obesity<sup>16</sup>. There has been a limited number of studies looking at the associations between obesity and diet type, but thus far results have been inconsistent and variable. Further work is therefore needed in order to determine the associations between diet type and obesity in cats and dogs</li> <li>Dogs prefer a higher fat and higher protein diet and, if this is fed <i>ad libitum</i>, dogs will overconsume and gain weight<sup>17</sup></li> </ul>
Reduced hair shedding	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Eliminated gas or bloat risk	<ul style="list-style-type: none"> <li>The suggested mechanism is that raw diets contain a decreased content of soluble and insoluble carbohydrate. While such a mechanism is plausible, there is currently no published evidence to support it</li> </ul>
Fewer food allergies and reduced skin complaints	<ul style="list-style-type: none"> <li>No scientific study evidence; anecdotal evidence only. Switching to a raw diet might plausibly resolve signs of an adverse reaction to food, provided that the component associated with the clinical signs is eliminated</li> </ul>
Reduced pancreatitis incidence	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Prevents parasites	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Lower veterinary costs overall	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Improved recovery from illness	<ul style="list-style-type: none"> <li>No scientific study evidence</li> <li>Nutrition should be tailored to the illness and a raw diet might not be appropriate for conditions such as renal disease, where phosphorus restriction is important<sup>18,19</sup></li> </ul>
Reduced body odour	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Reduced faecal odour	<ul style="list-style-type: none"> <li>Raw diets may reduce flatulence. This is thought to be due to having typically less fermentable fibre<sup>20</sup></li> <li>However, as a result of having less fermentable fibre, raw diets might result in decreased short-chain fatty acid production. This may be detrimental to the pet because short-chain fatty acids are essential for large intestinal health</li> </ul>
Increased energy levels	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Improved behaviour	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Improved immunity	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
The same diet can be fed at all life stages	<ul style="list-style-type: none"> <li>No scientific study evidence. Furthermore, some commercial producers of raw diets make different products for different life stages</li> <li>Homemade BARF-type diets are advertised as 'all-life' products. This implies that the recipe is suitable for young, old, pregnant, lactating, healthy or sick individuals. However, this is not necessarily advisable and any recipe should be formulated for the life stage of the pet, ideally by an appropriately qualified individual</li> </ul>
Fewer trips to the veterinary practice	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>
Increased mobility for older pets	<ul style="list-style-type: none"> <li>No scientific study evidence</li> </ul>

Overall, there is little evidence for many of the claims made, meaning further studies are required for veterinary professionals to deliver the best advice.

## What are the disadvantages of feeding raw diets?

### Risk of multidrug-resistant bacteria

- Associated risks are to:
  - The pet
  - The owners and their extended family – especially young, old and immunosuppressed individuals
  - Veterinary and hospital staff.
- Multidrug-resistant bacteria of particular concern include<sup>21</sup>:
  - Escherichia coli* (*E. coli*)
  - Meticillin-resistant *Staphylococcus pseudintermedius* (MRSP).



Delivered by BSAVA to:

Risk of pathogenic infection <sup>22</sup>

Pathogen type	Available evidence
<i>Salmonella</i> spp.	<ul style="list-style-type: none"> <li>Between 4% and 10% of chicken produced and sold in the UK for human consumption carries <i>Salmonella</i> spp. Amounts found on imported chicken are higher still <sup>23</sup></li> <li>Many studies have confirmed the transmission of <i>Salmonella</i> to pets in the contaminated meats that they are fed <sup>24,25,26,27,28</sup></li> <li>Simple cleaning of food bowls at home with soap and water or in a dishwasher is not sufficient to eliminate this pathogen</li> <li>Although uncommon, cases of zoonotic infection of <i>Salmonella</i> to pet owners have been linked to feeding a raw diet to the family pet <sup>29</sup></li> <li>Most pets that become infected with <i>Salmonella</i> become asymptomatic carriers, but studies have shown that they will still shed the bacteria in their saliva and faeces for 7 days after each infectious incident <sup>30</sup></li> </ul>
<i>E. coli</i>	<ul style="list-style-type: none"> <li>This pathogen was found in 60% of raw diets tested – both home-made and commercially produced <sup>31,32</sup></li> <li>Human deaths have been reported from handling raw dog food contaminated with <i>E. coli</i> <sup>22</sup></li> </ul>
<i>Clostridium</i> spp.	<ul style="list-style-type: none"> <li>Although the pathogenic nature of <i>Clostridium</i> spp. remains unclear, 20% of commercially produced raw diets were found to carry these bacteria <sup>33</sup></li> </ul>
<i>Campylobacter jejuni</i>	<ul style="list-style-type: none"> <li>This pathogen is present in 50–70% of all chicken sold for human consumption in the UK <sup>34</sup></li> <li>Acute polyradiculoneuritis in dogs (a neurological disorder) is thought to be triggered by pathogenic infection of <i>Campylobacter</i> deriving from contaminated food fed to pets <sup>35</sup></li> </ul>
<i>Toxoplasma gondii</i> spp.	<ul style="list-style-type: none"> <li>This pathogen is especially dangerous for pregnant women (as it causes fetal abnormalities) and for immunosuppressed individuals</li> <li>Women are advised to avoid direct contact with their cat's faeces and litter tray (as this is where the spread of the pathogen is most likely) during pregnancy <sup>36</sup></li> <li>As the prevalence of <i>Toxoplasma</i> in cats is greater when fed a raw diet, this should be given further consideration if living with at-risk individuals <sup>37,38,39</sup></li> </ul>
<i>Listeria</i> spp.	<ul style="list-style-type: none"> <li>This pathogen is widely found in meat products sold for human consumption <sup>40</sup>. Raw meat-based pet-food products have been withdrawn for containing dangerous levels of this pathogen <sup>41</sup></li> <li>Infection with <i>Listeria</i> spp. in humans causes a range of clinical signs from 'flu-like to fatal meningitis and abortion during pregnancy</li> </ul>
<i>Mycobacterium bovis</i>	<ul style="list-style-type: none"> <li>Found in uncooked commercially manufactured raw cat food <sup>42</sup></li> </ul>
Other pathogens found in raw diets include:	<ul style="list-style-type: none"> <li>All pose a health risk to all groups of people having contact with the pet</li> </ul> <ul style="list-style-type: none"> <li><i>Yersinia enterocolitica</i></li> <li>Tuberculosis</li> <li>Tularaemia</li> <li><i>Echinococcus multilocularis</i></li> </ul>

Concerns regarding the risks of pathogenic infection from raw diets have been frequently raised. A recent online survey taking a closer look at pet owners' perceptions of feeding raw diets and the incidence of infection, has suggested that the risks appear to be low, although not negligible <sup>43</sup>.



## Facts about pathogens

- Although not common, cases of pets suffering with *Salmonella* and other pathogens have been reported<sup>24,25,26,27,28</sup>.
- Risks to owners arise not only from the handling of uncooked food, but also from handling the pet – namely, contact with the pet's saliva, fur, urine and faeces are points of concern. Pathogens are shed by the pet during urination and defecation. During grooming, the pet may spread pathogens to their coat via their saliva.
- Pathogens may also be shed asymptotically into the environment where the pet lives or has contact with, which might be a further source of spread.
- 21–48% of home-prepared and commercially produced raw diets tested were found to have pathogenic contamination<sup>33,44</sup>.
- Freezing does not eliminate all pathogens that may be harmful. For example, it does decrease *Campylobacter* colony counts, but will not eradicate completely *Campylobacter* or other pathogens, some of which are unaffected by freezing. **The freezing of pet foods does not guarantee food safety**, as is commonly thought<sup>44</sup>.
- High hydrostatic pressure processing or high-pressure pasteurization is a process used by some commercial raw diet producers to reduce pathogen risk. However, it does not completely eradicate all pathogens and it is thought that it may lead to pathogenic resistance<sup>45</sup>.
- Some proponents of raw diets argue that the risk of pathogens from raw feeding is no greater than from any foods intended for human consumption. However, while there is undoubtedly a risk, it is arguably reduced because such food is usually cooked when consumed by humans. Furthermore, the risks associated with feeding raw food arise not only from the food but also from owner contact with the pet.
- Some proponents of raw feeding argue that contamination of pet food with pathogens is more common with conventional commercial foods, which are cooked. While this might be correct when examining the absolute number of pet food withdrawals, this is because such foods are much more commonly fed. Relatively speaking, many more come from raw-food manufacturers.
- **Based on available evidence, the risk of pathogenic infection should be considered when recommending a raw diet. Arguably, the potential risk to human health may be more of a concern than the risk to dogs and cats consuming raw diets. The prevalence of such infections is unknown, as it is not frequently reported, or in some cases the cause may not be associated with the pet or its food. Therefore, owners who wish to feed a raw diet should be strongly advised to take adequate precautions to reduce this risk.**

## Potential concerns from specific ingredients


Although known to be rare, there are some potential risks associated with specific types of food. It is important that owners wishing to feed a raw diet recognize these risks.

- Raw egg – contains avidin, which binds to the amino acid biotin making it unavailable. This may result in protein deficiency<sup>46</sup>.
- Raw fish (e.g. sea bream) or raw shellfish (e.g. scallops) – contains thiaminase, which destroys thiamine (an amino acid) causing protein deficiency<sup>46</sup>.
- Liver – fed in large quantities can cause hypervitaminosis A, due to the large amounts of vitamin A stored in liver tissue<sup>46</sup>.
- Trachea or gullet – if fed can cause hyperthyroidism, as the thyroid gland is often also consumed.

## Excess or deficiency

- When feeding a raw diet, especially if preparing it at home, it is very difficult for the pet owner to ensure that they are feeding a complete and balanced diet to their pet.
- Research looking to determine nutritional adequacy of home-prepared foods, both cooked and uncooked, has found some concerning results:
  - In one study where 95 different raw diets were examined<sup>47</sup>:
    - 60% had a significant imbalance
    - 40% had a minor imbalance or were balanced
  - In two similar studies examining 94 commonly available recipes for home-prepared diets (both cooked and uncooked) for pets with medical conditions all were found to be inadequate<sup>48,49</sup>
  - In a further study of 200 different recipes for maintenance diets, 95% were deficient in at least one essential nutrient, and 84% had multiple deficiencies<sup>50</sup>.
- Clinical signs associated with conditions that are caused by feeding an unbalanced diet can take a considerable period of time to appear. At this point the damage may be irreversible.
  - Deficiencies of certain nutrients include low calcium in the diet:
    - Occurs when an all-meat, table-scrap or all organ-meat diet is fed
    - These diets, being very low in calcium, cause hypocalcaemia resulting in nutritional secondary hyperparathyroidism. This is a very painful condition leading to significant bone loss through demineralization and multiple pathological bone fractures<sup>51,52</sup>.
- Although there are no published studies on prevalence, the risk of obvious nutrient deficiencies in adult dogs and cats appears to be rare. However, such deficiencies are more frequently reported in growing animals. As a result, caution should be exercised when feeding such diets to animals that have not yet reached skeletal maturity.

## Compliance

- As with any diet, compliance from owners will vary.
- As it is harder to balance a raw diet, the margin for error may be reduced.
- Arguably, the greatest concern comes from home-prepared raw diets. Not only are the recipes often unbalanced to begin with, but it is often difficult for owners to be fully compliant with the recipe. 

Delivered by BSAVA to:

## Can raw diets provide balanced nutrition?

Yes, provided that the recipe has been designed, overseen or approved by an appropriately qualified person.

## What safety measures are recommended to ensure that raw diets are fed responsibly?

### Excellent hygiene

Taking excellent hygiene precautions is essential for the safe feeding of raw diets. They should include the following:

- At purchase, inspect the ingredient packaging for damage or contamination. If damaged, the ingredient should not be purchased or should be discarded.
- Hand washing should commence:
  - After food preparation
  - After touching the pet.
- Preparation areas and surfaces should be cleaned with disinfectant immediately after food is prepared.
- Clean food bowls should be used for each meal.
- Bowls, floors and utensils should be disinfected immediately after feeding has finished.
- Bowls or plates for pet use should be kept separate from the ones used by the owners. Washing of bowls and plates alone is not enough to eliminate bacteria such as *Salmonella*<sup>53</sup>.
- Correct storage of foods will reduce the cross-contamination risk:
  - Any uncooked meats should be refrigerated in a sealed container
  - Uncooked meats should be stored away from human food (i.e. at the bottom of the refrigerator or, ideally, in a separate refrigerator – although this does not appear to be a contributing factor to the incidence of infection according to a recent study)<sup>43</sup>.
- Care is needed when handling pets and areas that the pet has contact with, especially areas used for elimination:
  - Studies have shown that pets infected with pathogenic bacteria can shed them into the environment<sup>22</sup>. Therefore, the following precautions should be observed:
    - Wear gloves or wash hands immediately after any handling or contact with urine or faeces
    - Swift disinfection of areas within the home where inappropriate elimination has occurred
    - Fastidious clearing or cleaning of elimination areas within the owner's garden
    - High general levels of environmental cleanliness and regular washing of bedding are also advisable, although further work is needed to ascertain whether washing would be sufficient for pathogen removal
    - Consideration should also be given to taking dogs to public areas, which may be difficult or impossible to clean (i.e. should these dogs be prevented from eliminating in public areas?).

### How to defrost frozen uncooked food safely

- Defrost uncooked foods in sealed containers or in the pet's bowl, ensuring that it is covered.
- Defrost uncooked foods on the bottom shelf of the refrigerator (ideally in a refrigerator that is separate from that storing food for human consumption).
- Do not refreeze thawed uncooked foods.
- Thaw juice will contain nutrients and should not be discarded but fed to the pet in the meal.
- Further storage of defrosted food must be in a sealed container and refrigerated.

### Method of feeding

Due to its high moisture content and potential for pathogenic contamination, it is not advisable to feed raw diets on an *ad libitum* basis. As with any food, an appropriate daily ration (based on the maintenance energy requirement of the dog or cat) should be fed in meals, with uneaten food being removed after 30 minutes, disposed of and never reoffered to the pet.

### Monitoring

It has been suggested that pets fed a raw diet should have regular tests to screen for nutrient deficiency. Tests should include biochemistry (including thyroxine concentrations), haematology and urinalysis. Adults should be tested every 6 to 12 months<sup>3</sup>.

## Considerations for feeding raw diets in healthy dogs and cats

- Dietary requirements can vary greatly between the different life stages and, arguably, no one diet will provide optimal nutrition for all stages.
- It is strongly advised to seek the advice of an appropriately qualified person before feeding a raw diet to growing animals. This is likely the life stage where optimal nutrition is most critical.

## Considerations for feeding raw diets in dogs and cats with various diseases

Many different diseases have specific nutritional requirements, meaning that a raw diet may not be appropriate in such circumstances. For example:

- **Chronic kidney disease (CKD):**
  - CKD requires a diet with restricted phosphorus content, while protein restriction should be considered at later stages<sup>54</sup>. This is extremely difficult to achieve with a raw diet, since protein and phosphorus content can be very high

Delivered by BSAVA to:

- **Obesity:**
  - Behavioural research indicates that dogs rarely regulate their food intake and will commonly overeat, especially when offered foods they find palatable<sup>55</sup>. The greater fat and protein content of raw diets may mean such diets are overconsumed if fed *ad libitum*, leading to unwanted weight gain. While some cats will regulate their own intake<sup>56,57</sup>, typically they consume more when fed higher protein diets
  - To avoid this, measured amounts of food, weighed carefully daily, should be fed
  - Regular monitoring of bodyweight is also essential to prevent obesity. Cats and dogs should be weighed every 2 weeks when transitioning to a new diet, with food intake adjusted until bodyweight is stable. They should then be weighed regularly thereafter (e.g. at least every 6 months to identify animals at risk of gradual weight gain)
- **Immune compromise:**
  - Pets that may be immunocompromised include those with neoplasia and hyperadrenocorticism, as well as those being treated with corticosteroids.

## Raw diets in a hospital environment – the rules

As more of our patients are being fed raw diets, it is important to consider the potential impact that this may have in a veterinary hospital. Many large hospitals and referral centres have adopted the following rules to reduce the risks to staff and other patients:

- No raw diets allowed into the hospital in order to:
  - Reduce risk to immunocompromised patients
  - Reduce cross-contamination risks to staff and other animals
- Patients consuming a raw diet at home may need to be isolated:
  - Pets can shed pathogens asymptotically for 7 days after eating contaminated meats.

## How to talk to owners about raw diets

### Remember:

- Most pet owners want to do what is best for their pet
- With the evidence available to us as veterinary professionals, and without prejudice, we must give the best advice that we can to ensure the pet receives a complete and balanced diet
- Veterinary professionals should separate what has been proven from advice we may be able to give based on clinical experience and reasoning
- When talking to owners about raw diets, it is important to understand their motivations, correct any misconceptions, and provide them with appropriate guidance so that they can feed their pet optimally and safely.

## Conclusion

The feeding of raw diets has become increasingly popular in recent years and it is therefore vitally important that we first educate ourselves on the available options, and then consider all the pros and cons when advising pet owners of the best diet to feed their pet. Decisions will often be made 'on balance' for the individual. Advice based on good evidence should be given each time and any recommendation must ensure that minimum requirements are met.

### References

1. Marx FR, Machado GS, Pezzali JG *et al.* (2016) Raw beef bones as chewing items to reduce dental calculus in Beagle dogs. *Australian Veterinary Journal* **94**(1–2), 18–23
2. Billinghurst I (1993) *Give your Dog a Bone: the Practical Common-Sense Way to Feed Dogs for a Long Healthy Life*. Bridge Printery, Alexandria NSW, Australia
3. Freeman LM, Chandler ML, Hamper BA and Weeth LP (2013) Current knowledge about the risks and benefits of raw meat-based diets for dogs and cats. *Journal of the American Veterinary Medical Association* **243**(11), 1549–1558
4. Hamper BA, Kirk CA and Bartges JW (2016) Apparent nutrient digestibility of two raw diets in domestic kittens. *Journal of Feline Medicine and Surgery* **18**(12), 991–996
5. Regulation (EU) 1334/2008 of the European Parliament and Council (2008) *On flavourings and certain food ingredients with flavouring properties for use in and on foods and amending Council Regulation (EEC) No. 1601/91, Regulations (EC) No. 2232/96 and (EC) No. 110/2008 and Directive 2000/13/EC*. [Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1585739671020&uri=CELEX:32008R1334>]
6. Axelsson E, Ratnakumar A, Arendt ML *et al.* (2013) The genomic signature of dog domestication reveals adaptation to a starch-rich diet. *Nature* **495**, 360–364
7. Morris GJ, Trudell J and Pencovic T (1977) Carbohydrate digestion by the domestic cat (*Felis catus*). *British Journal of Nutrition* **37**(3), 365–373
8. August JR, *et al.* (1985) Dietary hypersensitivity in dogs: cutaneous manifestations, diagnosis and management. *Compendium on Continuing Education for the Practising Veterinarian* **7**, 469–477
9. Harvey RG (1993) Food allergy and dietary intolerance in dogs: a report of 25 cases. *Journal of Small Animal Practice* **34**, 175–179
10. Hodgkins E (1991) Food allergy in cats: considerations, diagnosis and management. *Pet Vet* **24**, 8
11. Case LP, Carey DP, Hirakawa DA and Daristotle L (2000) *Canine and Feline Nutrition, 2nd edn*. Mosby Elsevier, Philadelphia
12. Vester BM, Burke SL, Liu KJ *et al.* (2010) Influence of feeding raw or extruded feline diets on nutrient digestibility and nitrogen metabolism of African wildcats (*Felis lybica*). *Zoo Biology* **29**, 676–686
13. Dogs Naturally (2020) *Raw Feeding Dogs: 10 Simple Rules to Get Started*. [Available from: [www.dogsnaturallymagazine.com/raw-feeding-primer](http://www.dogsnaturallymagazine.com/raw-feeding-primer)]

Delivered by BSAVA to:

14. Purina (2017) *What are Fillers in Dog Food?* [Available from: [www.justrightpetfood.com/blog/dog-food-fillers](http://www.justrightpetfood.com/blog/dog-food-fillers)]
15. Kendall PT and Holme DW (1982) Studies on digestibility of soya bean products, cereal, cereal and plant by-products in the diet of dogs. *Journal of the Science of Food and Agriculture* **33**, 813–820
16. German AJ, Woods GRT, Holden SL et al. (2018) Dangerous trends in pet obesity. *Veterinary Record* **182**, 25
17. Roberts MT, Bermingham EN, Cave NJ et al. (2018) Macronutrient intake of dogs, self-selecting diets varying in composition offered ad libitum. *Journal of Animal Physiology and Animal Nutrition* **102(2)**, 568–575
18. Finco DR, Brown SA, Crowell WA et al. (1992) Effects of dietary phosphorus and protein in dogs with chronic renal failure. *American Journal of Veterinary Research* **53(12)**, 2264–2271
19. Ross LA, Finco DR and Crowell WA (1982) Effect of phosphorus restriction on the kidneys of cats with reduced renal mass. *American Journal of Veterinary Research* **43(6)**, 1023–1026
20. Ugarte C, Guilford WG, Markwell P and Lupton E (2004) Carbohydrate malabsorption is a feature of feline inflammatory bowel disease but does not increase clinical gastrointestinal signs. *Journal of Nutrition* **8**, 2068S
21. Bos M, Broadfoot F, Healey K et al. (2019) *UK Veterinary Antimicrobial Resistance and Sales Surveillance*. [Available from: <https://www.gov.uk/government/publications/veterinary-antimicrobial-resistance-and-sales-surveillance-2018>]
22. Davies RH, Lawes JR and Wales AD (2019) Raw diets for dogs and cats: a review with particular reference to microbiological hazards. *Journal of Small Animal Practice* **60**, 329–339
23. Food Standards Agency (2003) *UK-wide Survey of Salmonella and Campylobacter contamination of fresh and frozen chicken on retail sale*. [Available from: [www.food.gov.uk](http://www.food.gov.uk)]
24. Cantor GH, Nelson Jr S, Vanek JA et al. (1997) *Salmonella* shedding in racing sled dogs. *Journal of Veterinary Diagnostic Investigation* **9(4)**, 447–448
25. Caraway CT, Scott AE, Roberts NC and Hauser GH (1959) Salmonellosis in sentry dogs. *Journal of the American Veterinary Medical Association* **135**, 599–602
26. Clark C, Cunningham J, Ahmed R et al. (2001) Characterization of *Salmonella* associated with pig ear dog treats in Canada. *Journal of Clinical Microbiology* **39(11)**, 3962–3968
27. Stone GG, Chengappa MM, Oberst RD et al. (1993) Application of polymerase chain reaction for the correlation of *Salmonella* serovars recovered from Greyhound faeces with their diet. *Journal of Veterinary Diagnostic Investigation* **5(3)**, 378–385
28. Joffe DJ and Schlesinger DP (2002) Preliminary assessment of the risk of *Salmonella* infection in dogs fed raw chicken diets. *Canadian Veterinary Journal* **43(6)**, 441–442
29. Minnesota Department of Health (2018) *Salmonella cases linked to raw meat dog food*. [Available from: [www.health.state.mn.us/news/pressrel/2018/salmonella020918](http://www.health.state.mn.us/news/pressrel/2018/salmonella020918)]
30. Verma AK, Sinha DK and Singh BR (2007) Salmonellosis in apparently healthy dogs. *Journal of Veterinary Public Health* **5(1)**, 37–39
31. Strohmeier RA, Morley PS, Hyatt DR et al. (2006) Evaluation of bacterial and protozoal contamination of commercially available raw meat diets for dogs. *Journal of the American Veterinary Medical Association* **228**, 537–542
32. Freeman LM and Michel KE (2001) Evaluation of raw food diets for dogs. *Journal of the American Veterinary Medical Association* **218**, 705–709 (Erratum published in *Journal of the American Veterinary Medical Association* 218, 1582)
33. Weese JS, Rousseau J and Arroyo L (2005) Bacteriological evaluation of commercial canine and feline raw diets. *Canadian Veterinary Journal* **46**, 513–516
34. Food Standards Agency (2016) *A UK wide microbiological survey of Campylobacter contamination in fresh whole chilled chickens at retail sale*. [Available from: [https://www.food.gov.uk/sites/default/files/media/document/retail\\_survey\\_protocol\\_year3\\_0.pdf](https://www.food.gov.uk/sites/default/files/media/document/retail_survey_protocol_year3_0.pdf)]
35. Martinez-Anton L, Marenda M, Firestone SM et al. (2018) Investigation of the role of *Campylobacter* infection in suspected acute polyradiculoneuritis in dogs. *Journal of Veterinary Internal Medicine* **32(1)**, 352–360
36. Gov.uk (2008) *Toxoplasmosis: diagnosis, epidemiology and prevention*. [Available from: [www.gov.uk/guidance/toxoplasmosis](http://www.gov.uk/guidance/toxoplasmosis)]
37. Smielewska-Los E, Rypula K and Pacon J (2002) The influence of feeding and maintenance system on the occurrence of *Toxoplasma gondii* infections in dogs. *Polish Journal of Veterinary Sciences* **5**, 231–234
38. Lopes AP, Cardoso L and Rodrigues M (2008) Serological survey of *Toxoplasma gondii* infection in domestic cats from north-eastern Portugal. *Veterinary Parasitology* **155**, 184–189
39. Dubey JP, Hill DE, Jones JL et al. (2005) Prevalence of viable *Toxoplasma gondii* in beef, chicken and pork from retail meat stores in the United States: risk assessment to consumers. *Journal of Parasitology* **91**, 1082–1093
40. Food Standards Agency (2019) *Listeria cases being investigated*. [Available from: <https://www.food.gov.uk/news-alerts/news/listeria-cases-being-investigated>]
41. Food Standards Agency (2019) *Raw Treat Pet Food Ltd recalls frozen raw beef, chicken, lamb, and chicken and tripe pet food due to the presence of Listeria monocytogenes*. [Available from: <https://www.food.gov.uk/news-alerts/alert/fsa-prin-36-2019>]
42. O'Halloran C, Ioannidi O, Reed N et al. (2019) Tuberculosis due to *Mycobacterium bovis* in pet cats associated with feeding a commercial raw diet. *Journal of Feline Medicine and Surgery* **21(8)**, 667–681
43. Anturaniemi J, Barrouin-Melo SM, Zaldivar-Lopez S et al. (2019) Owners' perception of acquiring infections through raw pet food: a comprehensive internet-based survey. *Veterinary Record* **185**, 658
44. Finley R, Reid-Smith R, Ribble C et al. (2008) The occurrence and antimicrobial susceptibility of Salmonellae isolated from commercially available canine raw food diets in three Canadian cities. *Zoonosis and Public Health* **55**, 462–469
45. Van Bree FPJ, Bokken GCAM, Mineur R et al. (2018) Zoonotic bacterial and parasites found in raw meat-based diets for cats and dogs. *Veterinary Record* **182(2)**, 50
46. Case LP, Carey DP, Hirakawa DA and Daristotle L (2000) *Canine and Feline Nutrition, 2nd edn*. Mosby Elsevier, Philadelphia
47. Dillitzer N, Becker N and Kienzle E (2011) Intake of minerals, trace elements and vitamins in bone and raw food rations in adult dogs. *British Journal of Nutrition* **106**, S53–S56
48. Larsen JA, Parks EM, Heinze CR and Fascetti AJ (2012) Evaluation of recipes for home-prepared diets for dogs and cats with chronic kidney disease. *Journal of the American Veterinary Medical Association* **240**, 532–538

Delivered by BSAVA to:



49. Heinze CR, Gomez FC and Freeman LM (2012) Assessment of commercial diets and recipes for home-prepared diets recommended for dogs with cancer. *Journal of the American Veterinary Medical Association* **241**, 1453–1460
50. Stockman J, Fascetti AJ, Kass PH et al. (2013) Evaluation of recipes of home-prepared maintenance diets for dogs. *Journal of the American Veterinary Medical Association* **242**, 1500–1505
51. Bennett D (1976) Nutrition and bone disease in the dog and cat. *Veterinary Record* **98**, 313–320
52. Hintz HF and Schryver HF (1987) Nutrition and bone development in dogs. *European Journal of Companion Animal Practice* **1**, 44–47
53. Weese JS and Rousseau J (2006) Survival of *Salmonella* Copenhagen in food bowls following contamination with experimentally inoculated raw meat: effects of time, cleaning and disinfection. *The Canadian Veterinary Journal* **47(9)**, 887–889
54. Polzin D (2019) *International Renal Interest Society—Diets for Cats with Chronic Kidney Disease (CKD)*. [Available from: [http://www.iris-kidney.com/education/protein\\_restriction\\_feline\\_ckd.html](http://www.iris-kidney.com/education/protein_restriction_feline_ckd.html)]
55. Roberts MT, Bermingham EN, Cave NJ et al. (2018) Macronutrient intake of dogs, self-selecting diets varying in composition offered *ad libitum*. *Journal of Animal Physiology and Animal Nutrition* **102**, 568–575
56. Beynen AC (2015) Dry foods for cats. *Creature Companion* **July**, 54–55
57. Beynen AC (2018) Cat food composition and caloric intake. *Dier-en-Arts Nr* **8/9**, 186–189

## Useful websites

The European Pet Food Industry

[www.fedialf.org](http://www.fedialf.org)

Food Standards Agency

- *Campylobacter*  
[www.food.gov.uk/safety-hygiene/campylobacter](http://www.food.gov.uk/safety-hygiene/campylobacter)
- *Listeria*  
[www.food.gov.uk/safety-hygiene/listeria](http://www.food.gov.uk/safety-hygiene/listeria)
- Pet Food – Business Guidance  
[www.food.gov.uk/business-guidance/pet-food](http://www.food.gov.uk/business-guidance/pet-food)
- *Salmonella*  
[www.food.gov.uk/safety-hygiene/salmonella](http://www.food.gov.uk/safety-hygiene/salmonella)

Gov.uk

- Raw pet foods: handling and preventing infection  
[www.gov.uk/guidance/raw-pet-foods-handling-and-preventing-infection](http://www.gov.uk/guidance/raw-pet-foods-handling-and-preventing-infection)

Pet Food Manufacturers' Association (PFMA)

[www.pfma.org.uk](http://www.pfma.org.uk)

World Small Animal Veterinary Association (WSAVA) – factsheets

[www.wsava.org](http://www.wsava.org)

- The Savvy Cat Owner's Guide: Nutrition on the Internet
- The Savvy Dog Owner's Guide: Nutrition on the Internet
- Selecting the Best Food for your Pet

## Last updated

April 2020